



Aigle, December 10, 2020

**Protocol for the resumption of  
BMX Race and BMX Freestyle events  
in the context of the new coronavirus pandemic**

**International BMX Race and BMX Freestyle events,  
including BMX Continental Championships,  
UCI BMX World Championships,  
UCI Urban World Championships and UCI BMX World Challenge.**

The stakeholders of cycling and the UCI acknowledge the extraordinary nature of the COVID-19 pandemic and the ensuing difficulties for the organisation of safe sporting events. These are all the more acute in the context of cycling events due to the regular international travel, the use of free-access venues and facilities and the different team and staff compositions.

In consideration of these extraordinary circumstances, it was agreed that the UCI proposes measures for the organisation of international BMX events (hereafter: the Recommendations). **These measures are mainly recommendations that apply to all international BMX Racing and BMX Freestyle events registered on the UCI calendar** (hereafter: the Events). The organiser has an obligation to carry out a risk assessment and to inform the stakeholders. The recommendations are not guidelines but should be seen as proposals for the organisation committees to reduce the risks of exposure and spread of the coronavirus that causes Covid-19.

In order to publish this protocol, the UCI relied on the one hand on the rules published for the organisation of international road cycling events (previously defined within an interdisciplinary steering committee) and on the specific risk analysis published by the World Health Organization (WHO), and reassessed by an international Task Force <sup>1</sup>

The protocol applies to all Events including BMX Continental Championships, UCI BMX World Cups, UCI BMX World Championships, UCI Urban World Championships and BMX World Challenge taking place as of approval by the UCI Management Committee until they are repealed by the UCI Management Committee and no earlier than 31 December 2021. It concerns all categories, men and women, elite, U23, juniors and BMX World Challenge competitors. The protocol is updated regularly taking into account new knowledge. Any modification will be published without delay and will be immediately applicable, unless otherwise indicated:

<https://www.uci.org/news/2020/covid-19-pandemic-how-to-return-to-cycling-events>

The document is divided into three main chapters,

- I. **List of mitigation measures for COVID-19**, a section setting out the practical recommendations to be implemented by organisers,
- II. **Risk assessment related to COVID-19**, a section concerning the risk assessment specifically related to COVID-19 (**mandatory measure**),
- III. **Risk assessment of the event**, a section defining the global risk assessment of the organisation of the Event (**mandatory measure**).

As a preamble, it is recalled that:

- local and national rules and laws prevail over the present protocol;
- the process of adapting the conditions for organising sporting events is part of a general risk-mitigation strategy, acknowledging however that the risks of infection may not be entirely excluded.

---

<sup>1</sup> Considerations for sports federations/sports event organizers when planning mass gatherings in the context of COVID-19. World Health Organization, 2020.

## I. List of mitigation measures for COVID-19

Specific risk mitigation measures are recommended to reduce the risk of transmission of the SARS-CoV-2 virus associated with the Events. It must be remembered that while mitigation measures may reduce the risk of infection with the novel coronavirus, they cannot completely eliminate the threat.

The list of recommended mitigation measures covers a wide range of subjects, including control of the risks of human-to-human transmission, emergency preparedness and response plans, coordination of actors and partners, communication, awareness campaigns on the means to fight and prevent COVID-19, etc.

The concrete actions to be implemented for an optimal organisation of BMX Race and BMX Freestyle competitions should be considered according to the national health regulations in force in the country (or administrative regions) of the Event, and according to the evaluation of the phase of the pandemic which will be made closer to the competition according to the criteria set out below (see paragraph II-B).

One of the globally acknowledged principles for organising cycling competitions is the creation and maintenance of protective "functional bubbles" around the riders and staff members. While this principle is applicable for all BMX Race events, it is not applicable for BMX Freestyle events. This discipline is an individual sport and generally riders do not stay in a team structure. Therefore, forming a "functional bubble" is unlikely.

For BMX Race events, the measures implemented should be based on the general objectives of

- controlling entry into functional bubbles by verifying the absence of virus carriage by suitable tests, **and**
- restricting direct and unprotected contact between members of these bubbles and third persons.

For BMX Freestyle events, the measures implemented should be based on the general objectives of

- verifying the absence of virus carriage by suitable tests, **and**
- restricting direct and unprotected contact with third persons by strictly applying individual protective measures (physical distancing, wearing a face mask, frequent hand washing).

In order to reduce the risks of spread and contamination by the new coronavirus, **the UCI recommends**, for the organisation of an Event, to apply the following measures:

### A- Pre-event measures

#### 1. Appointment of a COVID-19 Coordinator for the Event

An expert in infectious diseases should be appointed by the Event organiser; this COVID-19 Coordinator should have an up-to-date knowledge of the requirements and recommendations put in place by the national (or regional) health authorities to ensure the security of sporting events. He/she should get in touch with these authorities as soon as possible in order to best coordinate the actions to be implemented by the Event organiser with the rules in force. He/she regularly

consults the WHO website (<https://covid19.who.int>) or on a dedicated national website, to assess the pandemic status in the host country. This person is responsible for:

- determining the phase of the pandemic ahead of the competition, and is the advisor for the implementation of preventive measures. The COVID-19 Coordinator is the link between the Event organiser and the local or regional health authorities;
- assisting the Event organiser with the protocol for the management of suspected COVID-19 cases, including all stages of patient management until the diagnosis;
- providing the Event organiser the criteria for the identification of contact cases with a confirmed COVID-19 case (with either high-risk exposure, i.e. close contact, or low-risk exposure)<sup>2</sup> and coordinating the relevant actions with the local or regional health authorities.

## **2. Ensure that the accommodation where riders are staying is adequate to maintain a "life bubble"**

The UCI recommends that the organisers of BMX Race events offer accommodation arrangements enabling to maintain distancing between riders and staff members, and third persons. Measures such as grouping together on one floor (or a wing of the hotel), a reserved and independent dining room, help keep riders in a protection bubble. To ensure the application of the preventive measures put in place, the Event organiser is responsible for informing each hotel (e.g. room cleaning, physical distancing, hand washing, wearing a mask during service, etc.).

For BMX Freestyle events, the organisers must remind participants that their accommodation conditions must be compatible with the restriction of unprotected contact with third persons.

## **3. Ensure the isolation of suspected COVID-19 cases**

The UCI recommends that Event organisers reserve single rooms known as "isolation" to be used by anyone presenting symptoms suggestive of COVID-19, before referral to the COVID Doctor (see point I-B-3). We recommend booking 4 rooms for the duration of the event.

## **4. Inform the riders of the requirements and/or recommendations in terms of prevention procedures within their group (staff and riders)**

It is recommended that the organiser remind the importance of basic individual protection measures that must be taken by the riders and staff members in the official technical documents of the event. These measures include personal protection (physical distancing, wearing a mask), cleaning of technical equipment, cleaning and disinfection of commonly touched surfaces in the public areas, ventilation of confined areas, etc. **These measures should be appropriate to protect the integrity of the functional bubbles.** In this respect, the role of physicians and health professionals is essential.

---

<sup>2</sup> Information available on the websites of the European Centre for Disease Prevention and Control, and World Health Organisation.

## 5. Pre-Travel health checks

We recommend health checks for riders and staff members. The health checks are important before entering the protective bubbles (for BMX Race events), or before arriving on site (for BMX Freestyle events); they must include both a clinical and a biological component (both are complementary);

**a) the clinical aspect of detecting asymptomatic carriers** of the virus is based on examining clinical signs suggestive of the disease.

We recommend the use of a COVID clinical suspicion questionnaire, to be completed daily on the 5 days preceding the event. A questionnaire is **proposed below as a suggestion** (Figure 1). Like any medical questionnaire, it must be interpreted by a doctor, who may not be present on site. This is particularly important for riders in group or alone, without a doctor present on site. If it is used, adequate measures shall be taken in case the risk score is "highly suspicious " or "moderately suspicious" on 2 days out of 5. However, riders and staff members are free to use another clinical tool providing clinical guidance;


Covid-19 questionnaire	
Fever > 38°C	4 pts
Cough	4 pts
Shortness of breath	4 pts
Stuffy nose or sore throat	2 pts
Unusual aches	2 pts
Abnormal fatigue	2 pts
Unusual headache	1 pt
Diarrhea - vomiting	1 pt
<hr/>	
< or = 2	a little suspicious
3 - 5	moderately suspicious → PCR test according to the context
> or = 6	highly suspicious → PCR test

**Figure 1. Suggested screening questionnaire**

**b) the biological component** is based on the detection of the virus.

- the diagnosis of COVID-19 (biological component) is usually made using clinical, laboratory and radiological features. As symptoms and radiological findings of COVID-19 are non-specific, SARS-CoV-2 infection has to be confirmed by a molecular biology technique, mostly polymerase chain reaction (PCR), aimed at amplifying a specific RNA sequence of the SARS-CoV-2 virus. According to WHO, respiratory material for PCR should be collected from upper respiratory specimens (nasopharyngeal and oropharyngeal swabs or wash) in ambulatory patients. RT-PCR (Reverse Transcriptase-PCR) is a special PCR technique now being used to detect SARS-CoV-2. The new coronavirus can be detected in different tissues and body fluids and in clinical settings the respiratory material for PCR is mostly collected from naso-pharyngeal swabs.

- the general objective of the biological controls just before Events is to screen for healthy carriers of the virus or pre-symptomatic SARS-CoV-2 infections, and reduce the risk of transmission of the virus within the context of the Event. Specific procedures and tests need to be adapted to mass screening, while meeting a quality requirement with optimal sensitivity and specificity. Such screening tests intended for the qualitative detection of the SARS-CoV-2 RNA (i.e. viral tests) may be conducted as follows:
  - the use of saliva as an organic fluid for the detection of SARS-CoV-2<sup>3</sup>.
  - a highly specific and sensitive method based on the amplification of viral RNA<sup>4</sup>.
  - analyses on pooled individual salivary samples (pooling or multisampling methods)<sup>5</sup>.
- one qualitative test for the detection of SARS-CoV-2 RNA (PCR type) is recommended no more than 72 hours before the Event. If the organiser decides to have Covid tests carried out before the Event, the participation of a rider will only be authorized if the result of this test has been received before the Event and is confirmed as negative. (Figure 2 below).



**Figure 2. Schedule of pre-event PCR tests**

If the organisers decide to require a viral test (PCR type) before participating in the Event, they must set up a system to control the results of these tests respecting medical confidentiality and European data protection rules (GDPR).

## **6. Contacting the local health authorities (hospitals, emergency services)**

The UCI recommends that the COVID-19 coordinator of the event contact the local hospital and/or emergency medical services to inform them of the event, to ensure they have the capacity to handle trauma patients during the pandemic.

## **7. Ensuring all Event personnel have appropriate information on personal hygiene procedures;**

It is recommended that the organiser set up and ensure, as the event approaches, the strict application by the staff involved in the Event of individual measures to protect and prevent the spread of the virus.

<sup>3</sup> Saliva has been shown to be a viable alternative to nasopharyngeal swabs that cause discomfort due to procedure's invasiveness (Wyllie et al. 2020; Azzi et al. 2020)

<sup>4</sup> The technique used for the viral RNA identification must derive from PCR, such as RT-PCR, LAMP, RT-LAMP, SIBA, etc. (Jiang et al. 2020).

<sup>5</sup> (Lohse et al. 2020; Sunjaya et al. 2020). In order to preserve the sensitivity of the analyses it is recommended to constitute only pools lower or equal to 8 samples.

## **8. Arrange separate pathways for entry and exit**

- the BMX stadium
- all technical areas
- the media area
- the spectator areas to prevent congestion.

## **9. Arrange the communal areas accessible with accreditation to allow for physical distancing (min 1.5 m between people), especially;**

- in official areas
- in VIP areas.

Wearing of individual masks is highly recommended in these communal areas, especially when traveling within the enclosure. The obligation to wear a mask in VIP areas can only be lifted when sitting and drinking.

## **10. Provide hydro-alcoholic gel upon entrance and exit to the BMX stadium, technical zones, media centre and spectator areas.**

**B- During the Events,** the UCI recommends the following measures:

### **1. Identifying a physician in charge of COVID-19 suspected cases (COVID doctor)**

It is recommended to appoint a medical doctor responsible for managing any clinical suspicion of COVID-19 (“COVID doctor” for the Event), in coordination with local health services. Details on the management of suspected cases of COVID-19 are discussed later in paragraph I-B-2. The COVID doctor should have the mandatory protective equipment for medical personnel when dealing with COVID-19 suspected patients (FFP2 mask, gloves, visor or protective glasses, coveralls).

### **2. Management of a suspected COVID-19 case;**

- all persons involved in the Event (including riders and staff members) are requested to signal any suspicion of COVID-19 immediately to the Event medical services;
- the Event medical services will contact the COVID doctor to manage the suspect patient;
- the management of clinical cases will be carried out in agreement with the local or regional health services, and in accordance with WHO guidelines (see reference at the end of this document)
- the identification of contact cases with a confirmed COVID-19 case (close contacts and low-risk exposure contacts) will be the responsibility of the COVID doctor, in coordination with the competent health authorities;
- the implementation of the initial clinical examination protocol, and referral of the patient to the nearest COVID centre is the responsibility of the COVID doctor;
- it is recommended that the organisers make the details of these procedures available, as well as **the criteria for identifying risky contact cases, in the dedicated space provided by the UCI** (see chapter IV for the internet link).

### **3. Perform daily health checks** of riders during the Event;

- under the responsibility of the team (in general team doctor or physician remotely), or the rider himself/herself. It is at least recommended that the organiser remind the importance of the daily health checks in the official technical documents of the event.
- look for suspicious clinical signs using the questionnaire suggested above or other suitable tool;
- the check should be completed in the morning, before the daily activities.

### **4. Remind riders and staff members of the importance of wearing a mask** in all circumstances during the Event. Wearing a mask for riders and all staff members is compulsory on-site, except during training, warm-up sessions and competition for riders.

### **5. Organise the working conditions in the media centre.** Adapt the media centre reserved for the written and spoken press, both in terms of space, access and working conditions.

- organise the media centre to maintain a distance of 1.5 m between workstations, and provide hydro-alcoholic gel at the entrance and exit.
- the mixed zone will be enlarged, ventilated; all journalists should wear a mask, maintain a physical distance with the riders and use a pole mounted microphone for interviews.

### **6. Restrict spectators.**

- ban the presence of spectators in the finish area;
- limit spectators in the stands by assigning seating in order to provide physical distancing around each;
- wearing a mask for spectators should be compulsory inside the BMX stadium in all circumstances.

### **7. Limit access to the staging and start area.**

- limit access to the staging area to a maximum of 2 runs of 8 riders for BMX Racing, and 6 riders for BMX Freestyle;
- a distance of 1.5 m should be maintained between the lanes, and between 2 riders in the same lane;
- from the staging area to the starting ramp, riders should wear a mask;
- the masks will be placed in regulatory containers when entering the start gate.

### **8. Limit access to the finish area as much as possible.** Only allow access to the finish area for essential people (1 person per rider, a few photographers), and impose the wearing of a face mask).



**9. Create a cleaning schedule of any restrooms and disinfection of common areas and equipment;**

- regarding the toilets, ensure that there are enough stations on the site. Ensure the cleaning procedures that will be implemented, maintaining a physical distance of 1.5 m between users, including queues (to be respected using marks on the ground);
- regular cleaning of high touch areas and all contact points (door handles, switches, etc.);
- availability of hand sanitizers at strategic points.

**10. Provide waste bins for contaminated items** to allow for the safe disposal or storing of all hygienic materials.

**C- After the Event**

**1. Adjustment of the awards ceremony;**

The UCI recommends to:

- limit the size of the crowd, respecting social distancing (as per national health regulations)
- create 1.5 m pre-podium boxes in which riders can wait their turn to stand on the podium
- place the podium blocks 1.5 m apart
- require riders, and any other person involved, to wear a mask during the awards ceremony
- create a self-serve option where riders can collect their medals/trophies after hand sanitising
- request riders not to touch each other during the podium ceremony
- limit the number of photographers in front of the podium.

**2. Adapt the anti-doping station and procedures (compulsory measure);**

- ensure that doping control protocols are consistent with measures to prevent viral contamination (detection of asymptomatic carriers using viral tests (DCO, BCO) and chaperons, physical distancing outside and inside the station, procedures for checking and signing documents, etc.)
- a specific document is reported in Annex.

**II. Risk assessment related to COVID-19 (mandatory measure, included in the global risk assessment of the event, see chapter III)**

The first mandatory step with a view to organising an Event is for the Event organiser to carry out a preliminary risk assessment in accordance with national COVID-19 control strategies, if any. The aim of this risk assessment is to determine the overall risk of spreading the disease during the Event and the appropriate means to mitigate such a risk. This analysis is based on specific tools proposed by the World Health Organization (WHO), which have been revised and adapted by an International Task Force made of representatives from the world of sport.

risk of COVID-19 to the sporting event	Yes (1)/No (0)	Score
Will the event be held in a country that has documented active local transmission of COVID-19 (community spread)?	1	1
Will the event be held in multiple venues/cities/regions/countries?	1	1
Will the event include non-local/international participants (athletes and spectators) from areas that have documented active local transmission of COVID-19 (community spread)?	1	1
Will the event include a significant number of participants (athletes or spectators) at higher risk of severe COVID-19 disease (e.g., some athletes with disabilities, people with underlying health conditions)?	1	1
Will the event include conditions that could increase the risk of spread for COVID-19 (e.g. mass start or mass arrival, medical intervention, unavoidable contact or limited distancing measures)?	0	0
Will the event be held indoors?	0	0
<b>Total COVID-19 risk score</b>	<b>4</b>	<b>4</b>

**Figure 3. Specific COVID-19 risk score**  
(the numeric values are only given as examples)

The questions included in the COVID-19 risk assessment take into consideration the pandemic phase in the country of the Event, the risk factors linked to travel, human movement, and the possibility of the spread of the virus linked to characteristics of the competition itself


(Figure 3). The specific risks associated with indoor competitions on the spread of the virus are taken into account (item 6 of figure 3).

Completing this questionnaire gives a score that reflects the **specific risk associated with the pandemic**. The first question, which aims to characterize the state of the pandemic in the event region, deserves comment.

The different stages of a viral pandemic are clearly defined in a document published by the WHO, which describes the several phases of the influenza pandemic "*Pandemic influenza preparedness and response*". Although there are only few clinical and epidemiological analogies between the influenza and COVID-19 pandemics, the influenza transmission model is commonly used by health national agencies to characterise the stage of an epidemic. The different phases of an epidemic (which becomes a pandemic) can be illustrated according to the following diagram (Figure 4).

### A- The criteria

Different criteria are applied to characterise these phases, including both qualitative and quantitative factors. The difficulty is to propose criteria that are easily accessible in all countries of the world. The Event organisers should contact local or national health authorities in order to characterise the state of the pandemic according to the phases described by WHO. In order to make a first estimate, the following may be used as a basis:



**Figure 4. Current WHO pandemic phases**

- the number of new confirmed cases of COVID-19. The number of new cases reported each day is available for all countries in the world on the WHO website (<https://covid19.who.int>). Many websites regularly updated by national health authorities are now available. In order to smooth out the daily variations of figures, the weekly average may be considered. The daily number of new cases should be analysed for the country of the Event, and for other countries in the same WHO region.
- the basic reproductive number ( $R_0$ ) is an excellent parameter for characterising human-to-human transmission.  $R_0$  represents the number of people on average that a single infected individual may contaminate around him or her; it is a determining factor in epidemic risk assessment. A difficulty is obtaining this information for all countries. This information is not centralised by WHO and its estimation remains subject to the initiative of the national

authorities; the organisers should contact the national health authorities to obtain this information.

## **B- Characterisation of the different phases of the pandemic**

Although the decision of authorising a sporting event remains under the authority of the competent local or national authorities, the UCI considers that it can reasonably be considered that cycling events could be held during the following phases of the pandemic:

### **1- Community transmission of COVID-19 (WHO phases 5 and 6)**

This phase of the pandemic is characterized by confirmed human-to-human transmission of a coronavirus of animal origin, which can cause "epidemic outbreaks". Phases 5 and 6 mean that the pandemic state is imminent (WHO phase 5) or confirmed (WHO phase 6). It can be characterized by:

- confirmed cases in at least 2 countries in a WHO region and in at least 1 country outside the WHO region.
- a steady increase in the daily rate of confirmed clinical cases of COVID-19 (incidence rate). It is always difficult to interpret the data relating to new reported cases, as the modalities for diagnosing confirmed COVID cases depend on national strategies, either by systematic screening using RT-PCR (viral diagnostic) tests or by screening only contact cases and suspected patients of COVID-19 (also by RT-PCR tests), or only hospitalized patients, etc. This phase is characterized by clinical cases which appear in the form of extended clusters which progress towards generalized epidemic-type transmission.
- more than 50 new cases of COVID-19 declared per week, per 100,000 people;
- R0 values higher than 1.5.

### **2- Moderate risk period (WHO phase 4);**

This phase is characterised by confirmed human-to-human transmission of an animal-borne coronavirus, which can cause "outbreaks of epidemics". Phase 4 does not necessarily mean that a pandemic is inevitable. It can be characterised by:

- confirmed clinical cases occurring in only one country in a WHO region;
- a regular but moderate increase in the daily rate of confirmed clinical cases (difficult to quantify what is considered to be "moderate increase", since the methods of COVID-19 diagnosis depend on national strategies, either by systematic screening using RT-PCR tests for viral diagnosis, or by RT-PCR screening only of patients with suspected COVID-19 or having / who have been exposed to COVID-19, or only of hospitalised patients, etc. Furthermore, the data may not always be normalised to the global population). This phase is characterised by clinical cases present in the form of large clusters which tend to evolve towards a community transmission;
- 20 to 50 new cases of COVID-19 declared per week, per 100,000 people;
- R0 values higher than 1.5.

### **3- Low risk period (WHO phase 3, post-peak period);**

This low risk situation corresponds to either:

- the circulation of a coronavirus which causes sporadic infections or small clusters of respiratory infections. Human-to-human transmission does not appear to be sufficient to cause outbreaks. Limited human-to-human transmission can occur in certain circumstances of increased risk, but these modes of transmission remain limited to certain circumstances. This does not indicate that the virus has acquired the level of human transmissibility necessary to cause a pandemic. This period is a pandemic (pre-pandemic) alert period. This situation can be characterised by:
  - a sporadic and moderate increase in the daily rate of confirmed clinical cases.
  - R0 values higher than 1.5.
- the post-peak period of a pandemic. Pandemic activity appears to be decreasing but it is not certain whether or not new waves will occur. The drop in the level of activity of the pandemic should not mean the end of all preventive measures as several months may separate the arrival of new pandemic waves. This period can be characterised by:
  - a regular drop in the rate of confirmed COVID-19 cases. To assess this, the evolution of the average weekly reported COVID-19 cases can be monitored and be considered if there are less than 20 new cases declared per week per 100,000 people.
  - R0 values lower than 1.

### **4- Very low risk period (WHO phase 1, WHO phase 2, post-pandemic phase).**

This situation corresponds either to the identification of a coronavirus known to have caused infections in humans, and identified in wild and / or domestic animals (epizootic situation), or to a post-pandemic period during which the coronavirus will behave like a seasonal virus. At this stage, it is important to keep prevention measures to a minimum. We can characterise this period by:

- the absence of new confirmed cases for more than 3-4 weeks.
- R0 values less than 1 (for the post-pandemic period).

**The definition of the pandemic phase is the responsibility of the COVID-19 coordinator designated by the event organiser (see paragraph I-A-1).**

### III. Risk assessment of the event (mandatory measure)

The risk assessment is mandatory and allows organisers to review the main questions posed by the COVID-19 pandemic in the context of the organisation of an Event. This step helps the organisers understand and manage any specific risk associated with the pandemic.

This risk assessment should be reviewed regularly and updated immediately before the transition to the operational phase, depending on the risk mitigation measures in place, and in light of the evolution of the pandemic, which may be rapid. The organisers can refer to the guidelines and situation reports updated by the national public health authorities and / or the WHO (<https://covid19.who.int>).

#### It is carried out by combining,

- the analysis of the risks associated with COVID-19 (chapter II of the protocol), and
- the evaluation of risk mitigation measures.

The risk analysis is carried out using a dedicated Excel file available on the UCI website <https://www.uci.org/news/2020/covid-19-pandemic-how-to-return-to-cycling-events>

Part – BMX

File - "Risk-assess-BMX.xlsx".

#### A- Risks assessment related to COVID-19

The information from the questionnaire shown in Figure 3 (chapter II of the present protocol) should be reported on the sheet named "COVID" of the Excel file.

#### B- Risk mitigation measures.

Risk mitigation measures can be assessed using the sheet named "Measures" of the same Excel file. It includes each measure, each one being assigned a coefficient and the sum of the measures adopted determines the **risk mitigation score** that will be taken into account for the overall risk analysis of the event.

#### C- Matrix for the final decision.

It appears on the sheet named "Matrix" of the Excel file. The risk vs mitigation matrix combines the **COVID-19 specific risk score** and the **risk mitigation score** to determine a "colour" that identifies the total risk of transmission and spread of COVID-19 during the Event (Figure 5). This provides a clear indication of whether the staging of a sporting event is recommended or not, or whether other mitigation measures shall be required. The meanings of the colours are shown in the table below, with an overall risk determination.

The risk assessment should be repeated regularly, as soon as new preventive measures are implemented. The risk assessment and the defining of appropriate risk mitigation measures should, insofar as possible, be carried out with the involvement of local public health authorities and staff with expertise in mass gatherings, risk assessment, epidemiology and infectious disease control measures, from the very first stages of the Event planning.

### Overall risk score for the Event

The decision matrix takes the COVID-19 risk score and the mitigation score to provide a colour determination. This colour determination identifies the total risk of transmission and further spread of COVID-19 in relation to the mass gathering. The "Colour Determination" key below the decision matrix describes the total risk for each colour.

COVID-19 risk score	
Total mitigation score	

### COVID-19 risk Vs. Mitigation measures

		Total mitigation score			
		Very Prepared to Mitigate COVID-19 Impacts (76-100)	Somewhat Prepared to Mitigate COVID-19 Impacts (51-75)	Somewhat Unprepared to Mitigate COVID-19 Impacts (26-50)	Very Unprepared to Mitigate COVID-19 Impacts (0-25)
COVID-19 risk score	0 - Negligible	Very low	Very low	Very low	Very low
	1 - Very Low Risk	Very low	Very low	Low	Low
	2 - Low Risk	Low	Low	Low	Moderate
	3 - Moderate Risk (low-moderate)	Low	Moderate	Moderate	Moderate
	4 - Moderate Risk (high-moderate)	Moderate	Moderate	High	Very High
	5 - High Risk	High	High	Very High	Very High
	6 - Very High Risk	Very High	Very High	Very High	Very High

KEY FOR COLOUR DETERMINATION OF OVERALL RISK	
VERY LOW	Overall risk of transmission and further spread of COVID-19 in relation to the mass gathering is considered <b>very low</b> .
LOW	Overall risk of transmission and further spread of COVID-19 in relation to the mass gathering is considered <b>low</b> . Recommend checking whether mitigation measures can be strengthened.
MODERATE	Overall risk of transmission and further spread of COVID-19 in relation to the mass gathering is considered <b>moderate</b> . Recommend <b>significant</b> efforts to improve mitigation measures or reduce risk of transmission (decrease risk assessment score).
HIGH	Overall risk of transmission and further spread of COVID-19 in relation to the mass gathering is considered <b>high</b> . Recommend <b>significant</b> efforts to improve <b>both</b> mitigation measures and reduce risk of transmission (decrease risk assessment score).
VERY HIGH	Overall risk of transmission and further spread of COVID-19 in relation to the mass gathering is considered <b>very high</b> .

Figure 5. Total risk assessment score and interpretation

#### **IV- Exchange of information (mandatory measures)**

In order to promote the exchange of information necessary for the organisation of BMX Race and BMX Freestyle events, one secure data storage spaces will be opened by the UCI. This is intended for organisers to provide information to riders regarding the implementation of specific health-related measures.

**The link for this data storage space is as follows:**

<https://box.uci.ch/index.php/s/35GebIkTvcBHQ3p>

**The Event organisers shall deposit on this data storage space the 2 following documents, at the latest 2 weeks prior to the event:**

**A- the COVID-19 suspect case management protocol, including;**

- information concerning the phase of the pandemic in the region as the competition approaches, including incidence ratio (i.e. the number of Covid cases declared per week, per 100,000 persons, during the 2-3 weeks preceding the event)
- the procedures for managing suspected COVID-19 cases (i.e. availability of Covid laboratories recognized by the health authorities, operating availability, etc.)
- the criteria for defining contact cases, and their management.

**B- the result of the risk assessment,** using the file named “Risk-assess-BMX.xlsx”, and available on the UCI website (see chapter III). The risk assessment must include,

- the result of the risk assessment related to COVID-19 (sheet 1, “COVID”),
- a summary of the mitigation measures implemented (sheet 2, “Measures”),
- the overall risk score for the event (sheet 3, “Matrix”).

#### **V. Regulatory provisions**

Any subject or entity failing to implement the mandatory measures of the present protocol may be fined by the Disciplinary Commission between CHF 1,000 and CHF 10,000. The Disciplinary Commission shall determine the amount of the fine taking into account all the circumstances and in particular any aggravating or mitigating circumstances. Art. 12.2.005 of the UCI Regulations shall apply in case of a repeated offence.

Any subject or entity which defrauds, cheats or acts in an unfair manner when submitting the information required under this protocol to the UCI shall be sanctioned in accordance with article 12.4.008 of the UCI Regulations.



## References.

- Azzi L, Carcano G, Gianfagna F, Grossi P, Gasperina DD, Genoni A, Fasano M, Sessa F, Tettamanti L, Carinci F, Maurino V, Rossi A, Tagliabue A, Baj A. Saliva is a reliable tool to detect SARS-CoV-2. *J Infect.* 2020 Jul;81(1):e45-e50.
- Clinical management of COVID-19. Interim guidance. World Health Organization 2020, last version 27 May 2020.
- Considerations for sports federations/sports event organizers when planning mass gatherings in the context of COVID-19. World Health Organization 2020.
- Contact tracing: Public health management of persons, including healthcare workers, having had contact with COVID-19 cases in the European Union – first update. European Centre for Disease Prevention and Control. 31 March 2020
- Lohse S, Pfuhl T, Berkó-Göttel B, Rissland J, Geißler T, Gärtner B, Becker SL, Schneitler S, Smola S. Pooling of samples for testing for SARS-CoV-2 in asymptomatic people. *Lancet Infect Dis.* 2020 Apr 28.
- Pandemic influenza preparedness and response. A WHO guidance document. World Health Organization 2009. Reprinted 2010.
- Sunjaya AF, Sunjaya AP. Pooled Testing for Expanding COVID-19 Mass Surveillance. *Disaster Med Public Health Prep.* 2020 Jul 14:1-5.
- Wyllie A FJ, Casanovas-Massana A, Campbell M, Tokuyama M, et al. Saliva is more sensitive for SARS-CoV-2 detection in COVID-19 patients than nasopharyngeal swabs. <https://www.medrxiv.org/content/10.1101/2020.04.16.20067835v1> Web site. Published 2020. Accessed 12 May 2020.
- Yan C, Cui J, Huang L, Du B, Chen L, Xue G, Li S, Zhang W, Zhao L, Sun Y, Yao H, Li N, Zhao H, Feng Y, Liu S, Zhang Q, Liu D, Yuan J. Rapid and visual detection of 2019 novel coronavirus (SARS-CoV-2) by a reverse transcription loop-mediated isothermal amplification assay. *Clin Microbiol Infect.* 2020 Jun;26(6):773-779.

## ANNEX

# IN-COMPETITION TESTING SPECIFICITIES DURING COVID-19 BMX EVENTS

### MAKING HEALTH & SAFETY A TOP PRIORITY

December 2020

#### **1. CADF CONTROL OFFICER (DCO and BCO)**

When appointing a CADF Doping Control Officer (DCO) or Blood Collection Officer (BCO) for a BMX Race or BMX Freestyle event, the CADF has assessed that either is not at risk. SCP can be at risk if:

- they fall into a group of persons at risk; health care professionals working with COVID-19 positive patients, have tested athletes who tested positive to COVID-19 within a timeframe of 14 days after the mission, live with a person in one of the other risk groups or vulnerable populations.
- they fall into vulnerable persons' group due to age over 60 years' old, high blood pressure, diabetes, cardiovascular disease, compromised immune systems, etc., as advised by World Health Organization (WHO).

CADF DCOs & BCOs will perform a self-assessment (CADF document: cadf-034f\_rev0 - SCP self-assessment form) each day for the 5 days prior to the first planned controls. CADF DCOs & BCOs will have to have had a viral test, based on a PCR method, as far as possible 3 days before the first AD controls. Results will of course need to be negative for them to conduct the test. All documents will be submitted to the CADF & UCI using a dedicated online platform.

Based on the results, the CADF and the UCI Medical Director will decide whether to allow the CADF DCO or BCO to attend the event.

CADF DCOs and BCOs shall respect social distancing with any other sample collection or support personnel involved during the event.

#### **2. DOPING CONTROL STATION (DCS)**

A DCS must be provided by organisers as per UCI Testing & Investigations Regulations (UCI TIR).

In addition, organisers shall:

- ensure a spacious Doping Control Station (DCS) in order to ensure the recommended social distancing (at least 1.5 m) can be respected. Shouldn't the existing waiting room be spacious enough, please, consider setting an appropriate area for the athletes before the sample collection starts.
- provide premises that can be ventilated
- ensure the premises are cleaned and disinfected daily before use.
- provide disposable gloves. While gloves are not a substitute for hand hygiene, sample collection personnel (SCP) shall wear gloves throughout the sample collection process and athletes are also given the choice to wear gloves
- provide disposable face masks (medical face masks or non-medical masks or face covering); they shall be made available to the athlete, supporting personnel and SCP during the sample collection process.

- provide alcohol-based hand sanitizer
- provide disinfecting wipes and/or disinfecting spray
- provide disposable table cloth
- fence the area and provide someone to prevent non authorized persons to enter. Only one person is allowed to accompany the athlete.
- Provide waste bins for contaminated items to allow for the safe disposal or storing of all hygienic materials such as masks, gloves, etc.

### **3. DOPING CONTROLS IN HOTELS**

- Same prerequisites as listed above apply.
- Before conducting a doping control mission in a hotel, the DCO shall ensure that the tests can be conducted in a room that is ventilated and spacious enough to respect social distancing. If not possible, a minimum number of persons shall be present in the room; i.e. the athlete, the DCO, the BCO and if necessary, a physician.
- The SCP (DCO and chaperons) must regulate the arrival of athletes in the waiting room in the case where multiple athletes are tested. This will reduce the number of athletes in the same room.

### **4. NOTIFICATION PROCESS**

- Chaperons must be provided by organisers as usual according to UCI Testing & Investigations Regulations. Should the total risk of transmission and spread of COVID-19 be qualified by an Event organiser as higher than “moderate” in the total risk assessment (i.e. “community transmission”), as detailed in UCI’s procedure (see chapter II-B-1), chaperones should not be appointed. The assessment from the Covid coordinator will be available 2 weeks prior to the start of the race. On the day of the event, the chaperon will fill the self-assessment form.
- Chaperons will be responsible to notify athlete orally only respecting social distancing. A specific internal document for the chaperon will be created.
- The absence of signature of the rider and/or a third party upon oral notification does not prevent the rider to be bind.
- Should no chaperone be present, rider remains responsible for ensuring whether he/she has been selected to undergo Sample collection. The absence of a chaperone shall not excuse the rider for not reporting in time to the doping control station.
- List for notification purposes is displayed, where applicable usually near the finish line and near the DCS.
- It is the rider’s responsibility to remain within direct observation of the Chaperone at all times from the notification until the completion of the sample collection procedure.
- Rider must report immediately for sample collection and at the latest within 30 (thirty) minutes of finishing the Event, unless there are valid reasons for a delay, as per Article 7.4.2. of the UCI TIR.
- Written notification will be finalized with the DCO at the DCS
- In the event where the control would take place outside the DCS, such as in hotels (specific room or in rider’s/doctor’s room), as detailed before, only one athlete and one support personnel should be present at a time. When multiple riders are tested in hotels, notification will be done in a sensible manner but bearing in mind the no-advance notice aspect of these controls.

## 5. SAMPLE COLLECTION PROCESS

- In between athletes, surface where sample collection will take place must be cleaned using disinfectant wipes or disinfectant spray, including all materials to be used. As an alternative, a clean and disposable table cloth can be used.
- SCP must wash or sanitize hands and put on new gloves for each athlete and wear face mask.
- Athletes and supporting personnel (soigneur, doctor, etc.) must wear a face mask.
- Social/physical distancing is maintained as much as possible.
- Number of persons present during control session will be limited to minimum i.e.:
  - It is not necessary for organisers to provide a doctor/nurse to witness the miction, the task will be exceptionally ensured by the DCO if of the same gender. If not of the same gender, organisers will be asked to provide a doctor/nurse. On the day of the event, the doctor/nurse, if any, will fill the self-assessment form.
  - Only one person is allowed to accompany the athlete in the DCS area and during the sample collection process It is recommended that athletes present themselves at the DCS alone.

**NOTE:** Some specific situations may not allow the recommended distance to be maintained at all times. For example, **blood collection**, space limitations and/or the need for direct observation of urine sample provision are acceptable reasons to temporarily make allowances for closer distance.

## 6. COMPLETING SAMPLE COLLECTION SESSION

- Before leaving, work surfaces must be cleaned and all used materials (refractometer, pen, ruler etc.) cleaned with disinfectant wipes or spray.
- SCP must ensure that all discarded items/waste are disposed of in the appropriate bins for medical waste material.
- SCP guide athletes through the proper gloves and face mask removal techniques and ask them to place those items in their garbage bag.
- SCP instruct the athlete to clean their hands.

## 7. TRAMADOL CONTROLS SUPPORTED BY CADF

- Controls will be conducted in the Doping Control Station following the existing procedure at the end of events selected by the UCI, including the supplementary sanitary measures described above.
- The Tramadol Sample collection procedure may be amended if the circumstances so require.